

# MEASUREMENT WITHOUT ARCHIMEDEAN AXIOMS\*

LOUIS NARENS

*University of California, Irvine*

Axiomatizations of measurement systems usually require an axiom—called an *Archimedean axiom*—that allows quantities to be compared. This type of axiom has a different form from the other measurement axioms, and cannot—except in the most trivial cases—be empirically verified. In this paper, representation theorems for extensive measurement structures without Archimedean axioms are given. Such structures are represented in measurement spaces that are generalizations of the real number system. Furthermore, a precise description of “Archimedean axioms” is given and it is shown that in all interesting cases “Archimedean axioms” are independent of other measurement axioms.

**1. Preliminaries.** *Notation.* Throughout this paper the following convention will be observed.  $Re$  will stand for the real numbers;  $Re^+$  for the positive real numbers;  $I$  for the set of integers;  $I^+$  for the set of positive integers; and  $\langle x_1, \dots, x_n \rangle$  and  $(x_1, \dots, x_n)$  for ordered  $n$ -tuples; *iff* will stand for the phrase ‘if and only if’.

*Definition 1.1.* Let  $J$  be a nonempty set and  $\pi$  a function from  $J$  into the nonnegative integers. A *relational system of type  $\pi$*  is an ordered pair  $\langle A, \mathcal{F} \rangle$  where  $A$  is a nonempty set and  $\mathcal{F} = \{R_j \mid j \in J\}$  is a family of relations on  $A$  such that for each  $j \in J$ ,  $R_j$  is a  $\pi(j)$ -ary relation.

Comments on *Definition 1.1*:

- (1) By definition, a 0-ary relation on  $A$  is a member of  $A$ .
- (2) There is no bound on the cardinality of  $J$ . In fact, we will often use index sets  $J$  of cardinality greater than the continuum.
- (3) What is here called ‘relational systems’ are often elsewhere called ‘models (for first order languages)’.
- (4) Relations on  $A$  that have a special role are often listed separately from other relations. Thus if we are concerned with an ordering relation  $\lesssim$  on  $A$ ,  $\lesssim \in \mathcal{F}$ , we may write  $\langle A, \mathcal{F} \rangle$  as  $\langle A, \lesssim, \mathcal{F} \rangle$ , etc.
- (5) Operations on  $A$  can be represented as relations on  $A$ . For example, the two place operation  $+$  on  $A$  can be thought of as the three place relation  $R$  on  $A$  where  $R(a, b, c)$  holds if and only if  $a + b = c$ .

*Definition 1.2.* Let  $A$  be a nonempty set and  $\mathcal{F} = \{R_j \mid j \in J\}$  the set of all relations on  $A$ . Then  $\langle A, \mathcal{F} \rangle$  is called *the full relational system of  $A$  of type  $\pi$* , where  $\pi$  is the function from  $J$  into the nonnegative integers defined by:  $\pi(j) = n$  if and only if  $R_j$  is a  $n$ -ary relation.

*Definition 1.3.* Let  $J$  be a nonempty set and  $\mathcal{F} = \{R_j \mid j \in J\}$  and  $\pi$  a function from  $J$  into the nonnegative integers. By definition, *the language  $L_\pi(\mathcal{F})$*  is the first order language that has  $\mathcal{F} = \{R_j \mid j \in J\}$  as its set of predicate symbols where for each  $j \in J$ ,  $R_j$  is a  $\pi(j)$ -ary predicate symbol. (In  $L_\pi(\mathcal{F})$ ,  $\wedge$  is the conjunction symbol,  $\vee$  the disjunction symbol,  $\rightarrow$  the implication symbol,  $\leftrightarrow$  the “if and only if”

\* Received May, 1973.

symbol, and  $\neg$  the negation symbol.) If  $\langle A, \mathcal{G} \rangle$  is a relational system of type  $\pi$ , then by the usual definition of *truth* for first order languages (see [1], Chapter 3) each sentence of  $L_\pi(\mathcal{F})$  is assigned by  $\langle A, \mathcal{G} \rangle$  the truth value *true* or the truth value *false*.

Comments on *Definition 1.3*:

(1) 0-place predicate symbols are often called *individual constants* or *individual constant symbols*.

(2) Frequently, ' $\pi$ ' will be omitted from the expression ' $L_\pi(\mathcal{F})$ '. Thus, the expression ' $L(\mathcal{F})$ ' may be thought of as an abbreviation for ' $L_\pi(\mathcal{F})$ '.

*Definition 1.4.* Let  $\Gamma$  be a set of sentences of  $L_\pi(\mathcal{F})$  and  $\langle A, \mathcal{G} \rangle$  be a relational system of type  $\pi$ . Then  $\langle A, \mathcal{G} \rangle$  is said to be a *relational system for  $\Gamma$*  if and only if each sentence of  $\Gamma$  is true in  $\langle A, \mathcal{G} \rangle$ .

*Definition 1.5.* Let  $\langle A, \mathcal{F} \rangle$  and  $\langle B, \mathcal{G} \rangle$  be relational systems of type  $\pi$ . Then  $\langle A, \mathcal{F} \rangle$  and  $\langle B, \mathcal{G} \rangle$  are said to be *elementarily equivalent* if and only if each sentence of  $L_\pi(\mathcal{F})$  that is true in  $\langle A, \mathcal{F} \rangle$  is also true in  $\langle B, \mathcal{G} \rangle$ .

*Definition 1.6.* Let  $\Sigma$  be a set of sentences of  $L(\mathcal{F})$ .  $\Sigma$  is said to be *simultaneously satisfiable* if and only if there is a relational system  $\langle A, \mathcal{F} \rangle$  of type  $\pi$  such that each sentence of  $\Sigma$  is true in  $\langle A, \mathcal{F} \rangle$ .  $\Sigma$  is said to be *finitely satisfiable* if and only if each finite subset of  $\Sigma$  is simultaneously satisfiable. If  $\langle A, \mathcal{F} \rangle$  is a relational system such that each sentence of  $\Sigma$  is true in  $\langle A, \mathcal{F} \rangle$  then  $\langle A, \mathcal{F} \rangle$  is said to *simultaneously satisfy  $\Sigma$*  and  $\Sigma$  is said to be *simultaneously satisfiable in  $\langle A, \mathcal{F} \rangle$* .

The proof of the following fundamental theorem can be found in [1].

*Theorem 1.1. The Compactness Theorem of Logic.* If  $\Sigma$  is a set of sentences of  $L(\mathcal{F})$  that is finitely satisfiable then  $\Sigma$  is simultaneously satisfiable.

*Definition 1.7.* Let  $\langle A, \lesssim, \mathcal{F} \rangle$  be a relational system where  $\lesssim$  is a binary relation.  $\lesssim$  is said to be a *weak ordering on  $A$*  if and only if the following three sentences of  $L(\mathcal{F})$  are true in  $\langle A, \lesssim, \mathcal{F} \rangle$ :

- (1)  $\forall x(x \lesssim x)$ ,
- (2)  $\forall x \forall y \forall z((x \lesssim y \wedge y \lesssim z) \rightarrow x \lesssim z)$ ,
- (3)  $\forall x \forall y(x \lesssim y \vee y \lesssim x)$ .

*Definition 1.8.* If  $\lesssim$  is a weak ordering on  $A$  and  $a, b$  are elements of  $A$ , then, by definition,  $a \sim b$  iff  $a \lesssim b$  and  $b \lesssim a$ . It is easy to show that  $\sim$  is an equivalence relation on  $A$ .  $\lesssim$  is said to be a *total ordering on  $A$*  iff each equivalence class determined by  $\sim$  has exactly one member. By definition,  $a < b$  iff  $a \lesssim b$  and not  $a \sim b$ . Also by definition,  $a \gtrsim b$  iff  $b \lesssim a$ .

*Definition 1.9.* Let  $\langle A, \lesssim, \circ \rangle$  be a relational system where  $\lesssim$  is a binary relation and  $\circ$  a binary operation. Then  $\langle A, \lesssim, \circ \rangle$  is said to be an *ordered abelian group* if and only if it satisfies the following axioms:

- (1)  $\lesssim$  is a weak order on  $A$ ;
- (2) for each  $x$  and  $y$  in  $A$ ,  $x \circ y \sim y \circ x$ ;
- (3) for each  $x, y$ , and  $z$  in  $A$ ,  $x \circ (y \circ z) \sim (x \circ y) \circ z$ ;
- (4) for each  $x, y \in A$  there is a  $z \in A$  such that  $x \circ z \sim y$ ;
- (5) if  $x, y, z, w$  are in  $A$  and  $x \lesssim y$  and  $z \lesssim w$  then  $x \circ z \lesssim y \circ w$ .

Naturally, each of the axioms for an ordered abelian group can easily be formulated as a statement of  $L(\mathcal{F})$  where  $\mathcal{F} = \{\lesssim, \circ\}$ .

*Definition 1.10.* Let  $\langle A, \lesssim, \circ \rangle$  be an ordered abelian group. It is easy to show that there is a unique (up to equivalence) element  $e$  in  $A$  (called *the identity element*) such that  $a \circ e \sim a$  for all  $a$  in  $A$ . Let, by definition,  $A^+ = \{a \in A \mid e \prec a\}$ . By definition,  $|x|$  is a function from  $A$  into  $A$  such that (1) if  $e \lesssim a$  then  $|a| = a$  and (2) if  $e \succ a$  then  $|a| = b$  where  $b$  is some element of  $A$  such that  $a \circ b \sim e$ .

**2. Archimedean Axioms.** *Definition 2.1.* Let  $\langle A, \lesssim, \circ \rangle$  be an ordered abelian group. Let  $a \in A$ . Inductively define  $na$  for  $n \in I^+$  as follows:

- (i)  $1a = a$ ,
- (ii)  $(n + 1)a = (na) \circ a$ .

Let  $\mathcal{F} = \{\lesssim, \circ\}$ . In *Definition 2.1* note that ‘ $na$ ’ is not immediately formalizable as an expression of  $L(\mathcal{F})$  since ‘ $n$ ’ is not in the language of  $L(\mathcal{F})$ . But ‘ $3a$ ’ means ‘ $(a \circ a) \circ a$ ’ and this latter expression is easily formalizable in  $L(\mathcal{F})$ . In general, ‘ $na$ ’ is formalizable in  $L(\mathcal{F})$  for each  $n \in I^+$  by an expression that becomes increasingly longer for larger  $n$ .

Let  $\langle A, \mathcal{F} \rangle$  be a relational system of type  $\pi$ . In general, not all properties of  $\langle A, \mathcal{F} \rangle$  can be formulated in terms of  $L(\mathcal{F})$ . For example, if  $A$  is an infinite set, it can be shown that “the cardinality of  $A$ ” cannot be formulated in  $L(\mathcal{F})$ . In particular, “Archimedean” axioms cannot be formulated in  $L(\mathcal{F})$ .

*Definition 2.2.* Let  $\langle A, \lesssim, \circ \rangle$  be an ordered abelian group. Then  $\langle A, \lesssim, \circ \rangle$  is said to be *Archimedean* if and only if for each  $a, b$  in  $A^+$ , if  $a \prec b$  then for some  $n \in I^+$ ,  $b \lesssim na$ .

It is interesting to look at the parallel definition of ‘Archimedean’ using as much of the language  $L(\mathcal{F})$  as possible. Let  $\langle A, \lesssim, \circ, \mathcal{F} \rangle$  be the full relational system of  $A$  of type  $\pi$ . Assume that  $\langle A, \lesssim, \circ \rangle$  is an ordered abelian group. For each  $a \in A$  and each  $i \in I^+$  let the formula  $ia$  of  $L(\mathcal{F})$  be defined inductively as

$$1a = a,$$

$$(i + 1)a = ((ia) \circ a).$$

For each  $a, b$  in  $A$  and each  $i \in I^+$ , let  $\Psi_i(a, b)$  be the following sentence of  $L(\mathcal{F})$ :

$$\Psi_i(a, b): b \lesssim ia$$

Then  $\langle A, \circ, \lesssim \rangle$  is Archimedean if and only if for each  $a, b \in A^+$  there is an  $i \in I^+$  such that  $\Psi_i(a, b)$  is true in  $\langle A, \mathcal{F} \rangle$ . (Note that ‘for each  $a, b$  in  $A^+ \dots$ ’ can be formulated in  $L(\mathcal{F})$  by ‘ $\forall x \forall y \forall z (z \lesssim x \wedge z \lesssim y \wedge \forall x_1 (z \circ x_1 \sim x_1) \rightarrow \dots)$ ’). However, there is no way of formulating ‘there is an  $i \in I^+$  such that  $\Psi_i \dots$ ’ in  $L(\mathcal{F})$ .

In practice, Archimedean axioms take many different forms. We will now give a generalized definition of ‘Archimedean axiom’ that captures the essential quality of ‘Archimedeaness’.

*Definition 2.3.* Let  $\langle A, \mathcal{G} \rangle$  be a relational system and  $\langle A, \mathcal{F} \rangle$  a full relational system of  $A$ . (Thus  $\mathcal{G} \subseteq \mathcal{F}$ .) Let  $\mathcal{A} = \{\Psi_i(x_1, x_2, \dots, x_n) \mid i \in I^+\}$  be a set of formulas of  $L(\mathcal{F})$ .  $\mathcal{A}$  is said to be an *Archimedean schemata* for  $\langle A, \mathcal{G} \rangle$  if and only if

for each  $a_1, a_2, \dots, a_n$  in  $A$  there is an  $i \in I^+$  such that  $\Psi_i(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n)$  is true in  $\langle A, \mathcal{F} \rangle$ .

*Definition 2.4.* Let  $\Gamma$  be a set of sentences of  $L(\mathcal{G})$ . Let  $\mathcal{A} = \{\Psi_i(x_1, \dots, x_n) \mid i \in I^+\}$  be a set of formulas of  $L(\mathcal{G})$ . Then  $\mathcal{A}$  is said to be a *good Archimedean axiom* for  $\Gamma$  if and only if there is a relational system  $\langle A, \mathcal{G}' \rangle$  such that:

- (1)  $\langle A, \mathcal{G}' \rangle$  is a relational system for  $\Gamma$ , and
- (2)  $\mathcal{A}$  is an Archimedean schemata for  $\langle A, \mathcal{G}' \rangle$ .

The following theorem shows that in all interesting situations Archimedean axioms are not derivable from axioms expressible in  $L(\mathcal{G})$ .

*Theorem 2.1.* Let  $\Gamma$  be a set of sentences for  $L(\mathcal{G})$  and  $\mathcal{A} = \{\Psi_i(x_1, \dots, x_n) \mid i \in I^+\}$  be a good Archimedean axiom for  $\Gamma$ . Suppose that for each  $j \in I^+$  there is a relational system  $\langle A_j, \mathcal{G}_j \rangle$  of  $\Gamma$  such that there are elements  $a_1^j, a_2^j, \dots, a_n^j$  of  $A_j$  such that if  $\langle A_j, \mathcal{F}_j \rangle$  is the full relational system of  $A_j$  then for all  $i \leq j$ ,  $\neg \Psi_i(\mathbf{a}_1^j, \dots, \mathbf{a}_n^j)$  is true in  $\langle A_j, \mathcal{F}_j \rangle$ . Then there is a relational system  $\langle A, \mathcal{G}' \rangle$  of  $\Gamma$  for which the Archimedean axiom  $\mathcal{A}$  fails, i.e. there are  $a_1, \dots, a_n$  of  $A$  such that for each  $i \in I^+$   $\neg \Psi_i(\mathbf{a}_1, \dots, \mathbf{a}_n)$  is true in the full relational system of  $A$ .

*Proof.* Let  $\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n$  be new constant symbols that are not in the language  $L(\mathcal{G})$ . Let  $\Gamma' = \{\neg \Psi_i(\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n) \mid i \in I^+\}$ . We will show that  $\Sigma = \Gamma \cup \Gamma'$  is finitely satisfiable. Let  $\theta_1, \theta_2, \dots, \theta_n$  be finitely many statements of  $\Sigma$ . Without loss of generality suppose that  $\theta_1, \theta_2, \dots, \theta_k$  are members of  $\Gamma$  and  $\theta_{k+1}, \dots, \theta_n$  are members of  $\Gamma'$ . Since for  $k+1 \leq i \leq n$ ,  $\theta_i$  is  $\neg \Psi_{m_i}(\mathbf{b}_1, \dots, \mathbf{b}_n)$  for some  $m_i$ , let  $m$  be the maximum member of the set  $\{m_i \mid k+1 \leq i \leq n\}$ . By hypothesis, let  $\langle A_m, \mathcal{G}_m \rangle$  and  $a_1^m, \dots, a_n^m$  be such that

- (1)  $\langle A_m, \mathcal{G}_m \rangle$  is a relational system for  $\Gamma$ , and
- (2)  $a_1^m, \dots, a_n^m$  are in  $A_m$  and that for all  $i \leq m$

$\neg \Psi_i(\mathbf{a}_1^m, \dots, \mathbf{a}_n^m)$  is true in the full relational system of  $A_m$ . Interpret  $\mathbf{b}_1$  as  $a_1^m$ ,  $\mathbf{b}_2$  as  $a_2^m, \dots, \mathbf{b}_n$  as  $a_n^m$ . Then  $\theta_1, \theta_2, \dots, \theta_n$  are true in the full relational system of  $A_m$ . Thus  $\Sigma$  is finitely satisfiable. By the compactness theorem (*Theorem 1.1*), let  $\langle A, \mathcal{G}' \rangle$  be a relational system in which each sentence of  $\Sigma$  is true. Let  $a_1, \dots, a_n$  be the interpretations of  $\mathbf{b}_1, \dots, \mathbf{b}_n$  in  $\langle A, \mathcal{G}' \rangle$ . Then since for each  $i \in I^+$   $\neg \Psi_i(\mathbf{b}_1, \dots, \mathbf{b}_n)$  is true in  $\langle A, \mathcal{G}' \rangle$ ,  $\neg \Psi_i(\mathbf{a}_1, \dots, \mathbf{a}_n)$  is also true in  $\langle A, \mathcal{G}' \rangle$ . Thus  $\mathcal{A}$  fails in  $\langle A, \mathcal{G}' \rangle$ .

**3. Nonstandard Models of the Reals.** *Definition 3.1.* Let  $\langle Re, \mathcal{F} \rangle$  be the full relational system of  $Re$ .  $\langle *Re, *\mathcal{F} \rangle$  is said to be a *nonstandard model of the reals* if and only if the following three conditions hold:

- (1)  $\langle Re, \mathcal{F} \rangle$  and  $\langle *Re, *\mathcal{F} \rangle$  are elementarily equivalent in the language  $L(\mathcal{F})$ ;
- (2) if  $a \in Re$  then  $\mathbf{a}$  is interpreted in  $\langle *Re, *\mathcal{F} \rangle$  as  $a$ ; and
- (3)  $Re$  is a proper subset of  $*Re$ .

*Theorem 3.1.* There is a nonstandard model of the reals.

*Proof.* Let  $\langle Re, \mathcal{F} \rangle$  be the full relational system of  $Re$ . Let  $\Gamma$  be the set of sentences of  $L(\mathcal{F})$  that are true in  $\langle Re, \mathcal{F} \rangle$ . Let  $\mathbf{b}$  be a new constant symbol that is not in  $L(\mathcal{F})$ . Let  $\Gamma'$  be the following set of sentences:  $\Gamma' = \{\mathbf{b} \neq \mathbf{a} \mid a \in Re\}$ . Let  $\Sigma = \Gamma \cup \Gamma'$ . We will show that  $\Sigma$  is finitely satisfiable. Let  $\theta_1, \theta_2, \dots, \theta_n$  be members of  $\Sigma$ . Without loss of generality assume that  $\theta_1, \dots, \theta_k$  are in  $\Gamma$  and  $\theta_{k+1}, \dots, \theta_n$  are in  $\Gamma'$ . Then for each  $i, k+1 \leq i \leq n$ ,  $\theta_i$  is a sentence of the form  $\mathbf{a}_i \neq \mathbf{b}$  where  $\mathbf{a}_i \in Re$ . Since  $Re$  is an infinite set, let  $b' \in Re$  be such that  $b' \neq \mathbf{a}_i$  for each  $i, k+1 \leq i \leq n$ . Then  $\theta_1, \theta_2, \dots, \theta_n$  are true in  $\langle Re, \mathcal{F} \cup \{b'\} \rangle$  where  $\mathbf{b}$  is interpreted as  $b'$  and for each  $R \in \mathcal{F}$ ,  $\mathbf{R}$  is interpreted as  $R$ . By the compactness theorem (*Theorem 1.1*), let  $\langle *Re, *\mathcal{F} \cup \{b\} \rangle$  be a relational system of  $\Sigma$ . Let  $f$  be the following function from  $Re$  into  $*Re$ : for all  $a \in Re, f(a) = c$  where  $c$  is the interpretation of  $\mathbf{a}$  in  $*Re$ . Since each sentence of  $\Gamma$  is true in  $\langle *Re, *\mathcal{F} \rangle$ , it is easy to show that  $f$  is an isomorphic embedding of  $\langle Re, \mathcal{F} \rangle$  into  $\langle *Re, *\mathcal{F} \rangle$ . We may therefore assume that  $Re \subseteq *Re$  and for all  $a \in Re, a$  is the interpretation of  $\mathbf{a}$  in  $\langle *Re, *\mathcal{F} \rangle$ . Let  $b$  be the interpretation of  $\mathbf{b}$  in  $*Re$ . Since  $\mathbf{b} \neq \mathbf{a}$  is true in  $\langle *Re, *\mathcal{F} \cup \{b\} \rangle$  for each  $a \in Re$ , it follows that  $b \neq a$  for each  $a \in Re$ , i.e.  $b \in *Re - Re$ . Thus  $\langle *Re, *\mathcal{F} \rangle$  is a nonstandard model of the reals.

*Notation.* Let  $\langle Re, \mathcal{F} \rangle$  be the full relational system of  $Re$  and  $\langle *Re, *\mathcal{F} \rangle$  be a nonstandard model of the reals. Then for each  $n$ -place relation  $R$  in  $\mathcal{F}$  there is a predicate symbol  $\mathbf{R}$  in the language  $L(\mathcal{F})$ . In the relational system  $\langle Re, \mathcal{F} \rangle$ ,  $\mathbf{R}$  is interpreted as  $R$ . If  $R$  is a 0-ary relation, then it follows from the definition of nonstandard models of the reals that in  $\langle *Re, *\mathcal{F} \rangle$   $\mathbf{R}$  is interpreted as  $R$ . If  $R$  is a  $n$ -ary relation,  $n \geq 1$ , then, by convention, it will be assumed—unless otherwise explicitly stated—that the interpretation of  $\mathbf{R}$  in  $\langle *Re, *\mathcal{F} \rangle$  is  $*R$ .

Suppose that  $R$  is a  $n$ -place relation on  $Re$  where  $n \geq 1$  and that  $R(a_1, \dots, a_n)$ . Then  $\mathbf{R}(\mathbf{a}_1, \dots, \mathbf{a}_n)$  is true in  $\langle Re, \mathcal{F} \rangle$ . By elementary equivalence,  $\mathbf{R}(\mathbf{a}_1, \dots, \mathbf{a}_n)$  is also true in  $\langle *Re, *\mathcal{F} \rangle$ . Since in  $\langle *Re, *\mathcal{F} \rangle$   $\mathbf{a}_1, \dots, \mathbf{a}_n$  are interpreted as  $a_1, \dots, a_n$ , we can conclude that  $*R(a_1, \dots, a_n)$ . But this means that  $*R$  is an extension of  $R$ , i.e. that  $R \subseteq *R$ .

*Convention.* For convenience and clarity, the extensions of arithmetical operations and relations will often be denoted by the same symbol as those operations and relations of which they are extensions. That is,  $* <$  will often be written as  $<$ ,  $* =$  as  $=$ ,  $* +$  as  $+$ , etc.

*Definition 3.2.* Let  $\langle *Re, *\mathcal{F} \rangle$  be a nonstandard model of  $\langle Re, \mathcal{F} \rangle$ . An element  $\alpha$  in  $*Re$  is said to be *infinitesimal* if and only if for each  $r \in Re^+, |\alpha| < r$ .

*Theorem 3.2.* There is an infinitesimal  $\beta$  in  $*Re^+$ .

*Proof.* Since  $Re$  is a proper subset of  $*Re$ , let  $\alpha$  be in  $*Re$  and not be in  $Re$ . Since  $0 \in Re, \alpha \neq 0$ . Since  $\alpha \notin Re, -\alpha \notin Re$ . Since  $\forall x(x \neq 0 \rightarrow (0 < x \vee 0 < -x))$  is a true statement in  $L(\mathcal{F})$  of  $\langle Re, \mathcal{F} \rangle$  and therefore of  $\langle *Re, *\mathcal{F} \rangle$ , either  $0 < \alpha$  or  $0 < -\alpha$ . Without loss of generality, suppose that  $0 < \alpha$ .

Case 1.  $\alpha < r$  for each  $r \in Re^+$ . Then, by definition,  $\alpha$  is an infinitesimal and  $\alpha \in *Re^+$ .

Case 2.  $\alpha > r$  for each  $r \in Re^+$ . Let  $\beta = 1/\alpha$ . We will show that  $\beta$  is an infinitesimal and  $\beta \in *Re^+$ .  $\forall x(x > 0 \rightarrow 1/x > 0)$  is a true statement of  $\langle Re, \mathcal{F} \rangle$  and is therefore

a true statement of  $\langle *Re, *F \rangle$ . Since  $\beta > 0$  this means that  $1/\beta > 0$ . Hence  $\beta \in *Re^+$ . Let  $r$  be an arbitrary member of  $Re^+$ . Since  $1/r < \alpha$ , it is easy to show that  $\beta = 1/\alpha < r$ . But this means that  $\beta$  is an infinitesimal.

Case 3. There are  $r$  and  $s$  in  $Re^+$  such that  $r < \alpha < s$ . Let  $A_1 = \{t \in Re^+ \mid t < \alpha\}$  and  $A_2 = \{t \in Re^+ \mid t > \alpha\}$ . Then  $(A_1, A_2)$  forms a Dedekind cut of  $Re^+$ . Let  $c$  be the cut number determined by  $(A_1, A_2)$ . Since  $\alpha \notin Re^+$ ,  $\alpha \neq c$ . Therefore, either  $\alpha - c > 0$  or  $c - \alpha > 0$ . Without loss of generality, assume that  $c - \alpha > 0$ . (The case of  $\alpha - c > 0$  follows by a similar argument.) Let  $r$  be an arbitrary member of  $Re^+$ . Let  $\beta = c - \alpha$ . We will show that  $\beta < r$  thus establishing that  $\beta$  is an infinitesimal. Suppose that  $r \leq \beta$ . We will show a contradiction. Then  $r \leq c - \alpha$ , i.e.  $r + \alpha \leq c$ . Therefore,  $\alpha + r/2 < c$ . Thus  $c \in A_2$ . Let  $d = c - r/2$ . Then  $\alpha < d$ . Thus  $d \in A_2$ . Since  $d < c$ ,  $c$  cannot be the cut number of  $(A_1, A_2)$ ; a contradiction.

*Definition 3.3.* Let  $\langle *Re, *F \rangle$  be a nonstandard model of the reals and  $\alpha \in *Re$ .  $\alpha$  is said to be *finite* if and only if  $|\alpha| < r$  for some  $r \in Re$ .  $\alpha$  is said to be *infinite* if and only if  $|\alpha| > r$  for each  $r \in Re$ .

*Theorem 3.3.* If  $\alpha \in *Re$  and  $\alpha$  is finite then there is  $r \in Re$  such that  $\alpha - r$  is infinitesimal.

*Proof.* Let  $A_1 = \{t \in Re \mid t \leq \alpha\}$  and  $A_2 = \{t \in Re \mid t > \alpha\}$ . Then  $(A_1, A_2)$  forms a Dedekind cut of  $Re$ . Let  $r$  be the cut number of  $(A_1, A_2)$ . Then it is easy to show that  $\alpha - r$  is infinitesimal.

*Theorem 3.4.* If  $\alpha \in *Re$ ,  $s, r \in Re$ ,  $\alpha - s$  is infinitesimal, and  $\alpha - r$  is infinitesimal, then  $r = s$ .

Proof left to reader.

*Definition 3.4.* Let  $\langle *Re, *F \rangle$  be a nonstandard model of the reals and  $\alpha \in *Re$  and  $\alpha$  be finite. Then, by definition,  ${}^\circ\alpha$  is the unique  $r \in Re$  such that  $\alpha - r$  is infinitesimal.

*Theorem 3.5.* Let  $\alpha, \alpha_1$  be infinitesimal,  $\beta, \beta_1$  be finite, and  $\gamma, \gamma_1$  be infinite. Then the following are true:

- (1)  $\alpha + \alpha_1$  is infinitesimal,
- (2)  $\alpha\beta$  is infinitesimal,
- (3)  $\beta + \beta_1$  is finite,
- (4) if  $\beta$  and  $\beta_1$  are not infinitesimal, then  $\beta\beta_1$  is finite and not infinitesimal,
- (5)  $\beta + \gamma$  is infinite,
- (6) if  $\beta$  is not infinitesimal then  $\beta\gamma$  is infinite, and
- (7)  $\gamma\gamma_1$  is infinite

Proof left to reader.

**4. Imbeddings of Ordered Abelian Groups.** *Definition 4.1.* Let  $\langle A, \lesssim, \circ \rangle$  be an ordered abelian group. Let  $a, b$  be in  $A^+$ . Then  $a$  is said to be *commensurable with*  $b$  if and only if one of the following two conditions hold:

- (1)  $a \lesssim b$  and for some  $n \in I^+$   $b \lesssim na$ , or
- (2)  $b \lesssim a$  and for some  $n \in I^+$   $a \lesssim nb$ .

In the ordered abelian group  $\langle *Re, \leq, + \rangle$ , every pair of finite noninfinitesimal elements of  $*Re^+$  are commensurable. However, if  $\alpha$  is a positive infinitesimal, then  $\alpha$  and 1 are not commensurable.

*Theorem 4.1.* Let  $\langle A, \lesssim, \circ \rangle$  be an ordered abelian group. Then the relation defined by  $x$  being commensurable with  $y$  is an equivalence relation on  $A^+$ .

Proof left to reader.

*Definition 4.2.* The equivalence classes determined by the commensurability relation are called *commensurability classes*.

*Definition 4.3.* Let  $\langle A, \lesssim, \circ \rangle$  be an ordered abelian group. Then, by definition,  $\hat{A}$  is the set of commensurability classes of  $A^+$ . Also, if  $U, V$  are in  $\hat{A}$  then, by definition,  $U \lesssim V$  iff for some  $u \in U$  and some  $v \in V, u \lesssim v$ .

*Theorem 4.2.* Let  $A$  and  $\lesssim$  be as in *Definition 4.3*. Then  $\lesssim$  is a total ordering on  $\hat{A}$ .

Proof left to reader.

*Theorem 4.3.* Let  $\langle A, \lesssim, \circ \rangle$  be an Archimedean ordered group. Then  $\hat{A} = \{A^+\}$ .

Proof left to reader.

The following theorem shows that the commensurability classes need not be discretely ordered.

*Theorem 4.4.* Let  $A, B$  be commensurability classes of  $\langle *Re^+, \leq, + \rangle$  such that  $A < B$ . Then there is a commensurability class  $C$  such that  $A < C < B$ .

*Proof.* Let  $\alpha \in A$  and  $\beta \in B$ . Then for some  $\gamma \in *Re^+, \alpha\gamma = \beta$ . Since  $A < B, \alpha < \beta$ . Therefore  $\gamma > 1$ . Consider  $\delta = \alpha\sqrt{\gamma}$ . Since  $\gamma > 1, \alpha < \delta < \beta$ . Let  $n$  be an arbitrary member of  $I^+$ . Since  $A < B, n^2\alpha < \beta$ . Therefore  $n^2\alpha < \gamma\alpha = \beta$ . Thus  $n < \gamma = \sqrt{\gamma}$ . Therefore  $n\alpha < \sqrt{\gamma}\alpha = \delta$ . Since  $n$  is an arbitrary member of  $I^+$  we have shown that  $\delta$  is not in the commensurability class of  $\alpha$ , i.e.,  $\delta \notin A$ . Since  $n < \sqrt{\gamma}$  for each  $n \in I^+, n\delta < \sqrt{\gamma}\delta = \gamma\alpha = \beta$  for each  $n \in I^+$ . Thus  $\delta \notin B$ . Let  $C$  be the commensurability class that contains  $\delta$ . Then  $A < C$  and  $C < B$ .

Let  $\langle A, \lesssim, \circ \rangle$  be a relational system where  $\circ$  is a two place partial operation. Traditionally,  $\langle A, \lesssim, \circ \rangle$  is said to be an *empirical measurement system* if and only if there is a function  $G$  from  $A$  into  $Re^+$  that satisfies the following two conditions:

- (1) for each  $x, y$  in  $A, x < y$  iff  $G(x) < G(y)$ , and
- (2) for each  $x, y$  in  $A, G(x \circ y) = G(x) + G(y)$ .

Since the function  $G$  imbeds  $A$  into  $Re$  while preserving the intrinsic properties of  $\lesssim$  and  $\circ$ , many of the rich algebraic and topological properties of  $Re$  can be used for the analysis of the structure of  $\langle A, \lesssim, \circ \rangle$ . However, to guarantee that such a function  $G$  exists, it is necessary to assume (perhaps implicitly) some Archimedean axiom.

In the following it will be shown that ‘empirical measurement systems’ can be adequately axiomatized without use of Archimedean axioms. These measurement

systems will be imbedded in structures that are generalizations of the reals. Some of these,  $*Re$  for example, will have all the relational and algebraic properties of the reals.

*Definition 4.4.* Let  $\langle A, \lesssim, \circ \rangle$  be an ordered abelian group and  $S \subseteq A^+$ . Then a function  $f$  is said to be an *imbedding of  $S$*  (with respect to  $\lesssim, \circ$ ) into  $Re^+$  (respectively  $*Re^+$ ) if and only if the following three conditions hold:

- (1)  $f$  is a function from  $S$  into  $Re^+$  (respectively  $*Re^+$ );
- (2) for all  $x, y \in S, x < y$  iff  $f(x) < f(y)$ ; and
- (3) for all  $x, y \in S, f(x \circ y) = f(x) + f(y)$ .

*Theorem 4.5.* Let  $\langle A, \lesssim, \circ \rangle$  be an ordered abelian group,  $S \subseteq A^+$ , and  $f$  an imbedding of  $S$  into  $Re^+$  or  $*Re^+$ . Then the following two statements are true:

- (1) if  $x, y \in S$  and  $x \sim y$  then  $f(x) = f(y)$ , and
- (2) if  $x \in S$  and  $n \in I^+$  then  $f(nx) = nf(x)$ .

Proof left to reader.

*Theorem 4.6.* (Hölder's Theorem). Let  $\langle A, \lesssim, \circ \rangle$  be an Archimedean ordered abelian group. Then there is an imbedding of  $A^+$  into  $Re$ .

*Proof.* See [4], Chapter 2.

*Definition 4.5.* An ordered abelian group  $\langle A, \lesssim, \circ \rangle$  is said to be *regularly dense* if and only if for each  $x \in A$  and each  $n \in I^+$  there is a  $y \in A$  such that  $x \sim ny$ .

*Theorem 4.7.* (Abraham Robinson and Elias Zakon). Let  $\langle A, \lesssim, \circ \rangle$  be a regularly dense, ordered abelian group. Let  $\mathcal{F} = \{\lesssim, \circ\}$ . Then there is an Archimedean ordered abelian group that is elementarily equivalent to  $\langle A, \lesssim, \circ \rangle$  in the language  $L(\mathcal{F})$ .

*Proof.* See [13].

*Theorem 4.8.* Let  $\langle A, \lesssim, \circ \rangle$  be a regularly dense, ordered abelian group. Then there is a nonstandard model of the reals  $\langle *Re, *\mathcal{F} \rangle$ , an element  $u$  of  $*Re^+$ , and an imbedding  $G$  from  $A^+$  into  $*Re^+$  such that for each  $x$  in  $A, u < G(x)$ .

*Proof.* Let  $\mathcal{G} = \{\lesssim, \circ\}$ . By *Theorem 4.7*, let  $\langle B, \lesssim_1, \circ_1 \rangle$  be an Archimedean ordered group that is elementarily equivalent in the language  $L(\mathcal{G})$  to  $\langle A, \lesssim, \circ \rangle$ . Let  $D = B \cup Re$ . Let  $\langle D, \mathcal{H} \rangle$  be the full relational system of  $D$ . For notational simplicity, we will assume that  $D \cap A = \phi$ . Construct the first order language  $L_1$  from  $L(\mathcal{H})$  as follows: in each formula of  $L(\mathcal{H})$  replace each occurrence of  $\lesssim_1$  by  $\lesssim$  and each occurrence of  $\circ_1$  by  $\circ$ . Naturally, when interpreting the predicate symbols of  $L_1$  in  $\langle D, \mathcal{H} \rangle, \lesssim$  is interpreted as  $\lesssim_1$  and  $\circ$  as  $\circ_1$ . Let  $A_1 = \{a \mid a \in A\}$  be a new set of individual constant symbols and  $c$  still another new individual constant symbol. (In particular,  $c \notin A_1$ .) Let  $L_2$  be the first order language that has as its  $n$ -place predicate symbols ( $n \geq 1$ ) the  $n$ -place predicate symbols of  $L_1$ , and as its individual constant symbols the individual constant symbols of  $L_1$  together with  $A_1$  and  $\{c\}$ . Let  $\Gamma_0$  be



the set of sentences of  $L_1$  that are true in  $\langle D, \mathcal{H} \rangle$ . Let  $e$  be an identity element of  $\langle A, \lesssim, \circ \rangle$ . (Recall that the two place operation  $\circ$  is interpreted as a three place relation so that  $x \circ y = z$  stands for  $\circ(x, y, z)$ .) Let

$$\Gamma_1 = \{ \mathbf{x} \prec \mathbf{y} \mid x, y \in A \text{ and } x < y \} \cup \{ \mathbf{x} \circ \mathbf{y} = \mathbf{z} \mid x, y, z \in A \text{ and } x \circ y = z \}.$$

Let

$$\Gamma_2 = \{ \mathbf{c} \prec \mathbf{x} \mid x \in A^+ \} \cup \{ \mathbf{e} \prec \mathbf{c} \}.$$

Let

$$\Gamma = \Gamma_0 \cup \Gamma_1 \cup \Gamma_2.$$

We will now show that  $\Gamma$  is finitely satisfiable.

Let  $\Delta$  be a finite subset of  $\Gamma$ . Let  $\theta_1, \dots, \theta_m$  be the sentences of  $\Delta \cap (\Gamma_1 \cup \Gamma_2)$ . Let  $\mathbf{a}_1, \dots, \mathbf{a}_n$  be the individual constant symbols of  $A_1$  that occur in the formula  $\theta_1 \wedge \dots \wedge \theta_m$ . Let  $\Psi(\mathbf{c}, \mathbf{a}_1, \dots, \mathbf{a}_n)$  be the formula  $\theta_1 \wedge \dots \wedge \theta_m$ . Let  $a \in A^+$  be such that  $a < a_i$  for  $i = 1, \dots, n$ . (This can be done since  $\langle A, \lesssim, \circ \rangle$  is regular.) Then  $\Psi(\mathbf{a}, \mathbf{a}_1, \dots, \mathbf{a}_n)$  is true in  $\langle A, \lesssim, \circ \rangle$ . Therefore  $\exists x \exists x_1, \dots, \exists x_n \Psi(x, x_1, \dots, x_n)$  is true in  $\langle A, \lesssim, \circ \rangle$ . By elementary equivalence,  $\exists x \exists x_1, \dots, \exists x_n \Psi(x, x_1, \dots, x_n)$  is also true in  $\langle B, \lesssim_1, \circ_1 \rangle$ . Therefore let  $b, b_1, \dots, b_n$  be elements of  $B$  such that  $\Psi(b, b_1, \dots, b_n)$  is true in  $\langle D, \mathcal{H} \rangle$ . Then  $\Delta$  is simultaneously satisfiable in  $\langle D, \mathcal{H} \rangle$  under the following interpretation: each predicate symbol of  $\Delta \cap \Gamma_0$  is given its natural interpretation in  $\mathcal{H}$ ;  $\lesssim$  and  $\circ$  are interpreted as  $\lesssim_1$  and  $\circ_1$  respectively;  $\mathbf{a}_1, \dots, \mathbf{a}_n$  are interpreted as  $b_1, \dots, b_n$  respectively; and  $\mathbf{c}$  is interpreted as  $b$ .

By the compactness theorem (*Theorem 1.1*),  $\Gamma$  is simultaneously satisfiable in some relational system  $\mathcal{D}$ ,  $\mathcal{D} = \langle *D, * \lesssim, * \circ, A_2, * \mathcal{H} \rangle$ , where  $A_2$  is such that  $x \in A_2$  iff  $x$  is the interpretation of some  $y \in A_1$ . Let  $f$  be the following function from  $A^+$  onto  $A_2$ :  $f(x) = y$  iff  $y$  is the interpretation in  $\mathcal{D}$  of  $\mathbf{x}$ . Since for each  $x, y, z$  in  $A^+$ ,  $x \circ y = z$  iff  $\mathbf{x} \circ \mathbf{y} = \mathbf{z}$  is in  $\Gamma_1$  iff  $\mathbf{x} \circ \mathbf{y} = \mathbf{z}$  is true in  $\mathcal{D}$  iff  $f(x) * \circ f(y) = f(z)$ , and  $x < y$  iff  $\mathbf{x} \prec \mathbf{y}$  is in  $\Gamma_1$  iff  $\mathbf{x} \prec \mathbf{y}$  is true in  $\mathcal{D}$  iff  $f(x) * \prec f(y)$ , we can conclude that  $f$  is an isomorphic imbedding of  $A^+$  into  $A_2$ . Let  $c$  be the interpretation of  $\mathbf{c}$  in  $\mathcal{D}$ . Since  $\mathcal{D}$  is a relational system for  $\Gamma_2$ , for each  $x \in A_2$ ,  $c * \prec x$ . Let  $*B$  be the interpretation of  $\mathbf{B}$  in  $\mathcal{D}$ . Since  $\mathcal{D}$  is a relational system for  $\Gamma_1 \cup \Gamma_2$ ,  $A_2 \subseteq *B^+$  and  $c \in *B^+$ . Since  $\langle B, \lesssim_1, \circ_1 \rangle$  is an Archimedean ordered group, by *Theorem 4.6*, let  $F$  be an imbedding of  $\langle B, \lesssim_1, \circ_1 \rangle$  into  $\langle Re, \leq, + \rangle$ . Then  $F \in \mathcal{H}$ . Therefore the following sentences are in  $\Gamma_0$ :

- (1)  $\forall x(\mathbf{B}^+(x) \Leftrightarrow \exists y(\mathbf{Re}^+(y) \wedge \mathbf{F}(x) = y))$ ,
- (2)  $\forall x \forall y(\mathbf{B}^+(x) \wedge \mathbf{B}^+(y) \Rightarrow \mathbf{F}(x \circ y) = \mathbf{F}(x) + \mathbf{F}(y))$ ,
- (3)  $\forall x \forall y((\mathbf{B}^+(x) \wedge \mathbf{B}^+(y)) \Rightarrow (x \prec y \Leftrightarrow \mathbf{F}(x) < \mathbf{F}(y)))$ .

Since  $\mathcal{D}$  is a relational system for  $\Gamma_0$ , the above three sentences are true in  $\mathcal{D}$ . Let  $*F$  be the interpretation of  $\mathbf{F}$  in  $\mathcal{D}$ . Then,

- (1') for each  $x \in *B^+$  there is a  $y \in *Re^+$  such that  $*F(x) = y$ ;
- (2') for each  $x, y$  in  $*B^+$ ,  $*F(x \circ y) = *F(x) + *F(y)$ ; and
- (3') for each  $x, y$  in  $*B^+$ ,  $x * \prec y$  iff  $*F(x) < *F(y)$ .

Recall that  $f$  is an imbedding of  $\langle A^+, \lesssim, \circ \rangle$  into  $\langle *B^+, *\lesssim, *\circ \rangle$ . For each  $x \in A^+$  let  $G(x) = *F(f(x))$ . Then  $G$  is an imbedding of  $\langle A^+, \lesssim, \circ \rangle$  into  $\langle *Re^+, \lesssim, \circ \rangle$ . Since  $c * \prec x$  for each  $x \in A_2$ ,  $*F(c) < G(x)$  for each  $x \in A^+$ .

*Definition 4.6.* Let  $\langle A, \lesssim, \circ \rangle$  be an ordered abelian group and  $S \subseteq A^+$ . Then  $B$  is said to be a *set of units for S* if and only if the following three conditions hold:

- (1)  $B \subseteq S$ ;
- (2) for each  $x \in S$  there is a  $y \in B$  such that  $y$  is commensurable with  $x$ ; and
- (3) if  $x$  and  $y$  are in  $B$  and  $x \neq y$  then  $x$  and  $y$  are not commensurable.

*Theorem 4.9.* Let  $\langle A, \lesssim, \circ \rangle$  be an ordered abelian group and  $S \subseteq A^+$ . Then there is a set of units for  $S$ .

Proof left to reader.

*Definition 4.7.* Let  $\langle A, \lesssim, \circ \rangle$  be an ordered abelian group and  $S \subseteq A^+$ . A *scale s for S* is a function from  $S$  into  $Re^+$  such that the following five conditions hold:

- (1) if  $x, y \in S$  and  $x$  is commensurable with  $y$  then  $s(x \circ y) = s(x) + s(y)$ ;
- (2) if  $x, y \in S$  and  $x$  is commensurable with  $y$  then: (i) if  $x \lesssim y$  then  $s(x) \leq s(y)$ , and (ii) if  $s(x) < s(y)$  then  $x \prec y$ ;
- (3) if  $x, y \in S$ ,  $x \prec y$ , and  $x$  is not commensurable with  $y$ , then  $s(x \circ y) = s(y)$ ;
- (4) if  $x, y \in S$ ,  $x$  is commensurable with  $y$ ,  $z \in A^+$ ,  $x \circ z \sim y$ , and  $s(x) = s(y)$ , then  $z$  is not commensurable with  $x$ ; and
- (5)  $B = \{x \mid x \in S \text{ and } s(x) = 1\}$  is a set of units for  $S$ .

$B$  is called the *set of units for s*.

*Theorem 4.10.* Let  $\langle A, \lesssim, \circ \rangle$  be a regularly dense ordered abelian group,  $S \subseteq A^+$ , and  $B$  a set of units for  $S$ . Then there is a scale  $s$  for  $S$  such that  $B$  is the set of units for  $s$ .

*Proof.* Let  $\langle Re, \mathcal{F} \rangle$  be the full relational system for  $Re$ . By *Theorem 4.8* let  $\langle *Re, *\mathcal{F} \rangle$ ,  $\alpha$  and  $G$  be such that the following four conditions hold:

- (1')  $\langle *Re, *\mathcal{F} \rangle$  is an elementary extension of  $\langle Re, \mathcal{F} \rangle$ ;
- (2')  $\alpha \in *Re^+$ ;
- (3')  $G$  is an imbedding of  $A^+$  into  $*Re^+$ ; and
- (4') for each  $x \in A^+$ ,  $G(x) > \alpha$ .

The following sentence of  $L(\mathcal{F})$  is true in  $\langle Re, \mathcal{F} \rangle$ :

$$\forall x \forall y ((\mathbf{Re}^+(x) \wedge \mathbf{Re}^+(y) \wedge x \prec y) \rightarrow \exists z (\mathbf{I}^+(z) \wedge zx \leq y \wedge (z + 1)x > y)).$$

Therefore, by elementary equivalence, for each  $x, y$  in  $*Re^+$ , if  $x \prec y$  then for some  $z$  in  $*I^+$ ,  $zx \leq y$  and  $(z + 1)x > y$ . Therefore, for each  $x \in S$ , let  $N_x$  be a member of  $*I^+$  such that  $N_x \alpha \leq G(x)$  and  $(N_x + 1)\alpha > G(x)$ . Let  $a, b$  be members of  $S$ . We will show that the following two statements are equivalent:

- (i)  $a$  and  $b$  are commensurable,
- (ii)  $N_a/N_b$  is finite and not infinitesimal.

Without loss of generality, suppose that  $a \lesssim b$ . Suppose (i). Let  $n \in I^+$  be such that  $na \lesssim b$  and  $(n+1)a \succ b$ . Then

$$N_b \alpha \leq G(b) < G((n+1)a) = (n+1)G(a) < (n+1)(N_a + 1)\alpha.$$

Thus  $N_b \alpha < (n+1)(N_a + 1)\alpha$ . Dividing by  $\alpha$  we get  $N_b < (n+1)(N_a + 1)$ . Thus

$$\frac{1}{n+1} \leq \frac{N_a + 1}{N_b} \leq \frac{2N_a}{N_b}.$$

That is,

$$\frac{1}{2(n+1)} < \frac{N_a}{N_b}.$$

Thus we have shown that  $N_a/N_b$  is not infinitesimal. Since  $a \lesssim b$ , it follows that  $N_a \leq N_b$ , i.e.  $N_a/N_b \leq 1$ , i.e.  $N_a/N_b$  is finite. We have therefore shown that (i) implies (ii). Now suppose (ii). Since  $N_a/N_b$  is not infinitesimal, let  $q \in I^+$  be such that  $1/q < N_a/N_b$ . Then  $(N_b + 1) < (q + 1)N_a$ . Therefore

$$G(b) < (N_b + 1)\alpha < (q + 1)N_a\alpha \leq (q + 1)G(a) = G((q + 1)a).$$

Therefore  $b < (q + 1)a$ . That is,  $a$  is com measurable with  $b$ . Therefore (ii) implies (i).

We need the following two lemmas to complete the proof:

*Lemma 1:* if  $x \in A^+$  then  $N_x$  is infinite.

*Proof:* Let  $x \in A^+$  and  $n$  be an arbitrary member of  $I^+$ . Since  $\langle A, \lesssim, \circ \rangle$  is regularly dense, let  $z \in A^+$  be such that  $nz \sim x$ . Since  $z \in A^+$ ,  $\alpha < G(z)$ . Therefore  $n\alpha < nG(z) = G(nz) = G(x)$ . That is, for each  $n \in I^+$ ,  $n\alpha < G(x)$ . Therefore for each  $n \in I^+$ ,  $n < N_x + 1$ , i.e.  $N_x$  is infinite.

*Lemma 2:* if  $x, y \in A^+$  then  $N_x + N_y - 1 < N_{x \circ y} < N_x + N_y + 2$ .

*Proof:*

$$N_x \alpha + N_y \alpha \leq G(x) + G(y) = G(x \circ y) < (N_{x \circ y} + 1)\alpha$$

and

$$N_{x \circ y} \alpha \leq G(x \circ y) = G(x) + G(y) < (N_x + 1)\alpha + (N_y + 1)\alpha = (N_x + N_y + 2)\alpha.$$

For each  $x \in S$  let  $\beta(x)$  be the unit of  $B$  that is com measurable with  $x$ . For each  $x \in S$  let

$$s(x) = \circ \left( \frac{Nx}{N_{\beta(x)}} \right).$$

We now show that  $s$  is a scale for  $S$ . Since  $\beta(x)$  is com measurable with  $x$ ,  $N_x/N_{\beta(x)}$  is finite but not infinitesimal. Therefore,

$$\circ \left( \frac{N_x}{N_{\beta(x)}} \right) \in Re^+$$

for each  $x \in S$ . That is,  $s$  is a function from  $S$  into  $Re^+$ .

(1) Suppose  $x, y \in S$  and  $x$  is commensurable with  $y$ . Then  $x \circ y$  is commensurable with  $x$ . Therefore  $\beta(x) = \beta(y) = \beta(x \circ y)$ . Let  $N = N_{\beta(x)}$ . Then by *Lemma 1*,  $N$  is infinite. By *Lemma 2*,

$$\frac{N_x + N_y - 1}{N} < \frac{N_{x \circ y}}{N} < \frac{N_x + N_y + 2}{N}.$$

Since  $N$  is infinite,  $1/N$  and  $2/N$  are infinitesimal. Therefore by simple algebra

$$\begin{aligned} s(x) + s(y) &= \circ\left(\frac{N_x + N_y}{N}\right) = \circ\left(\frac{N_x + N_y - 1}{N}\right) \leq \circ\left(\frac{N_{x \circ y}}{N}\right) = s(x \circ y) \\ &\leq \circ\left(\frac{N_x + N_y + 2}{N}\right) = \circ\left(\frac{N_x + N_y}{N}\right) = s(x) + s(y). \end{aligned}$$

That is,  $s(x) + s(y) \leq s(x \circ y) \leq s(x) + s(y)$ , i.e.  $s(x \circ y) = s(x) + s(y)$ .

(2) Let  $x, y \in S$  be such that  $x$  is commensurable with  $y$ . Then  $\beta(x) = \beta(y)$ . Let  $N = N_{\beta(x)}$ . Then, (i) if  $x \lesssim y$  then  $N_x \leq N_y$  and  $\circ(N_x/N) \leq \circ(N_y/N)$ , i.e.  $s(x) \leq s(y)$ , and (ii) if  $s(x) < s(y)$  then  $\circ(N_x/N) < \circ(N_y/N)$ , i.e.  $N_x < N_y$ , i.e.  $x < y$ .

(3) Let  $x, y \in S$  be such that  $x \lesssim y$  and  $x$  is not commensurable with  $y$ . Since  $y < x + y < y + y$ ,  $y$  is commensurable with  $x + y$ . Therefore  $\beta(y) = \beta(x \circ y)$ . Let  $z = \beta(y)$ . Since  $x$  is not commensurable with  $y$  and  $y$  is commensurable with  $z$ ,  $x$  is not commensurable with  $z$ . Since  $x < y$ , it then follows that  $x < z$ . Therefore, for each  $n \in I^+$ ,  $nx < z$ . Thus for each  $n \in I^+$ ,  $G(nx) = nG(x) < G(z)$ . That is,  $nN_x\alpha < (N_z + 1)\alpha$  for each  $n \in I^+$ . That is, for each  $n \in I^+$ ,  $nN_x - 1 < N_z$ . Thus

$$\frac{N_x}{N_z} < \frac{1}{n} + \frac{1}{nN_z} < \frac{1}{n-1}$$

for each  $n \in I^+$ ,  $n \geq 2$ . By *Definition 3.2*, this means that  $N_x/N_z$  is infinitesimal. Since by (1),  $s(x \circ y) = s(x) + s(y)$ , we can conclude that

$$s(x \circ y) = s(x) + s(y) = \circ\left(\frac{N_x + N_y}{N_z}\right) = \circ\left(\frac{N_y}{N_z}\right) = s(y).$$

(4) Let  $x, y \in S$  and  $z \in A^+$  be such that  $x$  is commensurable with  $y$ ,  $x \circ z \sim y$ , and  $s(x) = s(y)$ . Then we will show by contradiction that  $z$  is not commensurable with  $x$ . Assume that  $z$  is commensurable with  $x$ . Let  $n \in I^+$  be such that  $x \lesssim nz$ . Then  $N_x\alpha < n(N_z + 1)\alpha$ . That is,  $N_x < n(N_z + 1) = nN_z + n$ . Thus,

$$\frac{N_x - n}{n} = \frac{1}{n}N_x - 1 < N_z.$$

Since  $x$  is commensurable with  $y$ ,  $\beta(x) = \beta(y)$ . Let  $N = N_{\beta(x)}$ . Then, by *Lemma 2*,

$$N_x + \left(\frac{1}{n}N_x - 1\right) - 1 < N_x + N_z - 1 < N_{x \circ z} = N_y.$$

Since, by *Lemma 1*,  $1/N$  is infinitesimal and

$$\circ\left(\frac{N_y}{N}\right) = s(y) = s(x) = \circ\left(\frac{N_x}{N}\right),$$

we can conclude that

$$\left(1 + \frac{1}{n}\right) \circ \left(\frac{N_x}{N}\right) = \circ \left(\frac{N_x + \left(\frac{1}{n}N_x - 1\right) - 1}{N}\right) \leq \circ \left(\frac{N_y}{N}\right) = \circ \left(\frac{N_x}{N}\right).$$

That is,

$$\left(1 + \frac{1}{n}\right) \circ \left(\frac{N_x}{N}\right) \leq \circ \left(\frac{N_x}{N}\right)$$

which is impossible since

$$\circ \left(\frac{N_x}{N}\right) \in Re^+.$$

*Theorem 4.11.* Let  $\langle A, \lesssim, \circ \rangle$  be an ordered abelian group,  $S \subseteq A^+$ , and  $s, t$  be scales for  $S$ . Let  $T$  be a subset of  $S$  that satisfies the following two conditions:

- (1) for each  $x, y \in T$ ,  $x$  is commensurable with  $y$ ; and
- (2) for each  $x \in T$ ,  $nx \in T$  for sufficiently large positive integers  $n$ .

Then there is a positive real number  $r$  such that for all  $x \in S$ ,  $s(x) = rt(x)$ .

*Proof.* Let  $x \in T$  and  $r = s(x)/t(x)$ . Let  $y \in T$ . Then for each sufficiently large  $n \in I^+$ , let  $N_n \in I^+$  be such that  $N_n y \lesssim nx < (N_n + 1)y$ . Then  $N_n$  approaches infinity as  $n$  approaches infinity. Also  $s(N_n y) \leq s(nx) \leq s((N_n + 1)y)$ . From which we conclude,  $N_n s(y) \leq ns(x) \leq (N_n + 1)s(y)$ . Therefore

$$\frac{N_n}{n} \leq \frac{s(x)}{s(y)} \leq \frac{N_n + 1}{n}.$$

Similarly,

$$\frac{N_n}{n} \leq \frac{t(x)}{t(y)} \leq \frac{N_n + 1}{n}.$$

Letting  $n$  approach infinity, we have

$$\lim \frac{N_n}{n} = \frac{s(x)}{s(y)} = \frac{t(x)}{t(y)}.$$

Since  $s(x) = rt(x)$ , we get

$$\frac{rt(x)}{s(y)} = \frac{t(x)}{t(y)},$$

i.e.  $rt(y) = s(y)$ .

**5. Extensive Structures.** *Definition 5.1.* Let  $\langle A, \lesssim, \circ \rangle$  be a relational system such that  $\lesssim$  is a two place relation on  $A$ ,  $B \subseteq A \times A$ , and  $\circ$  is a function from  $B$  into  $A$ . (I.e.  $\circ$  is a partial operation on  $A$ .) Then  $\langle A, \lesssim, \circ \rangle$  is said to be an *extensive structure* if and only if the following six conditions are satisfied for all  $a, b, c \in A$ :

- (1)  $\lesssim$  is a weak order on  $A$ ;

- (2) if  $(a, b) \in B$  and  $(a \circ b, c) \in B$ , then  $(b, c) \in B$ ,  $(a, b \circ c) \in B$ , and  $(a \circ b) \circ c \succsim a \circ (b \circ c)$ ;
- (3) if  $(a, c) \in B$  and  $a \succsim b$ , then  $(c, b) \in B$  and  $a \circ c \succsim c \circ b$ ;
- (4) if  $a \succ b$ , then there exists  $d \in A$  such that  $(b, d) \in B$  and  $a \succsim b \circ d$ ;
- (5) if  $(a, b) \in B$ , then  $a \circ b \succ a$ ; and
- (6) there exist  $x, y \in A$  such that  $a \prec x \circ y$ .

*Definition 5.2.* The relational system  $\langle A, \succsim, \circ \rangle$  is said to be an *Archimedean extensive structure* if and only if the following two conditions hold:

- (1)  $\langle A, \succsim, \circ \rangle$  is an extensive structure; and
- (2) if  $a_1, a_2, \dots, a_n, \dots$ , is an (infinite) sequence of members of  $A$  such that for  $n = 2, 3, \dots, a_n = a_{n-1} \circ a_1$ , then for all  $b \in A$  it is not the case that for each  $n \in I^+$ ,  $a_n \prec b$ .

*Theorem 5.1.* Let  $\langle A, \succsim, \circ \rangle$  be an Archimedean extensive structure and  $B$  be the domain of  $\circ$ . Then there is a function  $f$  from  $A$  into  $Re^+$  such that the following two conditions hold for all  $x, y \in A$ :

- (1)  $x \prec y$  iff  $f(x) < f(y)$ ; and
- (2) if  $(x, y) \in B$  then  $f(x \circ y) = f(x) + f(y)$ .

Furthermore, if  $g$  is another function from  $A$  into  $Re^+$  that satisfies (1) and (2), then there is  $r \in Re^+$  such that for all  $z \in A$ ,  $g(z) = rf(z)$ .

*Proof.* See [4], Chapter 3.

*Theorem 5.2.* Let  $\langle A, \succsim, \circ \rangle$  be an ordered abelian group,  $S \subseteq A^+$ , and  $s, t$  scales for  $S$ . Let  $a \in S$  and  $T = \{y \in S \mid y \text{ is commensurable with } a\}$ . Suppose that  $T$  satisfies the following three conditions:

- (1) if  $x, y, z \in T$ ,  $x \circ y \in T$ , and  $z \prec y$ , then  $x \circ z \in T$ ;
- (2) if  $x, y \in T$  and  $x \prec y$ , then there is a  $z \in T$  such that  $x \circ z \in T$  and  $x \prec x \circ z \prec y$ ; and
- (3) if  $x \in T$  then there are  $y, z \in T$  such that  $x \prec y \circ z$ .

Then there is a positive real number  $r$  such that for all  $x \in T$ ,  $s(x) = rt(x)$ .

*Proof.* Let  $\succsim_1$  and  $\circ_1$  be the restrictions of  $\succsim$  and  $\circ$  to  $T$ . Then it is easy to verify that  $\langle T, \succsim_1, \circ_1 \rangle$  is an extensive structure. Since all members of  $T$  are commensurable,  $\langle T, \succsim_1, \circ_1 \rangle$  is Archimedean. Therefore, by *Theorem 5.1*, there is a positive real number  $r$  such that for all  $x \in T$ ,  $s(x) = rt(x)$ .

*Definition 5.3.* The relational system  $\langle A, \succsim, \circ \rangle$  is said to be a *closed extensive structure* if and only if the following five conditions hold for all  $a, b, c \in A$ :

- (1)  $\succsim$  is a weaker order on  $A$ ;
- (2)  $\circ$  is a binary operation on  $A$ ;
- (3)  $(a \circ b) \circ c \succsim a \circ (b \circ c)$ ;
- (4) if  $a \succsim b$  then  $a \circ c \succsim c \circ b$ ; and
- (5)  $a \circ b \succ a$ .

*Theorem 5.3.* Let  $\langle A, \lesssim, \circ \rangle$  be a closed extensive structure. Then the following are true for all  $a, b, c \in A$ :

- (1) if  $a \gtrsim c$  and  $b \gtrsim d$  then  $a \circ b \gtrsim c \circ d$ ;
- (2)  $a \circ b \sim b \circ a$ ;
- (3)  $a \circ (b \circ c) \sim (a \circ b) \circ c$ .

Proof left to reader.

*Theorem 5.4.* Let  $\langle A, \lesssim, \circ \rangle$  be a closed extensive structure. Let

$$D = \{(x, y) \mid x, y \in A\},$$

$$P = \{(x, y) \mid x, y \in A \text{ and } x \gtrsim y\},$$

$$\oplus \text{ be the 2-place operation on } D \text{ such that } (a, b) \oplus (c, d) = (a \circ c, b \circ d),$$

$$(a, b)^{-1} = (b, a), \text{ and}$$

$${}_1 \gtrsim \text{ be such that } (a, b) {}_1 \gtrsim (c, d) \text{ iff } (a, b) \oplus (c, d) \in P.$$

Then  $\langle D, \lesssim_1, \oplus \rangle$  is an ordered abelian group. Furthermore,  $\langle A, \lesssim, \circ \rangle$  is isomorphic to a subset of  $D^+$ . (That is, there is a one-to-one function  $f$ ,  $f(x) = (2x, x)$ , from  $A$  into  $D^+$  such that (1)  $x \lesssim y$  iff  $f(x) \lesssim_1 f(y)$  and (2)  $f(x \circ y) = f(x) \oplus f(y)$ .)

Proof left to reader.

*Theorem 5.5.* Every ordered abelian group is a subgroup of a regularly dense ordered abelian group.

*Proof.* Let  $\langle A, \lesssim, \circ \rangle$  be an ordered abelian group. Let  $e$  be an identity element of  $A$ . For each  $x \in A$ , let  $x^{-1}$  be an element of  $A$  such that  $x \circ x^{-1} \sim e$ . For each  $n \in I^+$  and each  $x \in A$ , define  $(-n)x$  to be  $nx^{-1}$ . Let  $B = \{(x, n) \mid x \in A \text{ and } n \in I - \{0\}\}$ . Define the two place operation  $\oplus$  on  $B$  as follows:

$$(x, n) \oplus (y, m) = ((mx) \circ (ny), nm).$$

Define the two place relation  $\lesssim_1$  on  $B$  as follows:  $(x, n) \lesssim_1 (y, m)$  if and only if one of the following two conditions hold:

- (1)  $nm$  is positive and  $e \lesssim ny \circ (-m)x$ , or
- (2)  $nm$  is negative and  $e \gtrsim ny \circ (-m)x$ .

It is easy to verify that  $\langle B, \lesssim_1, \oplus \rangle$  is an ordered abelian group. It is easy to show that for each  $n \in I^+$  and each  $(x, m) \in B$  that  $n(x, nm) = (x, m)$ . Thus  $\langle B, \lesssim, \oplus \rangle$  is regularly dense. One can also verify that the function  $f$ , defined by  $f(x) = (x, 1)$  iff  $x \in A$ , is an isomorphic imbedding of  $\langle A, \lesssim, \circ \rangle$  into  $\langle B, \lesssim_1, \oplus \rangle$ . We may therefore consider  $\langle A, \lesssim, \circ \rangle$  as a subgroup of  $\langle B, \lesssim_1, \oplus \rangle$ .

*Definition 5.4.* Let  $\langle A, \lesssim_1, \circ_1 \rangle$  be a relational system where  $\lesssim_1$  is a binary relation on  $A$  and  $\circ_1$  is a binary partial operation on  $A$  (i.e.  $\circ_1$  is a function from  $D \times D$  into  $A$  for some  $D \subseteq A$ ). Then  $\langle B, \lesssim, \circ \rangle$  is said to be a *regularly dense ordered abelian group extension* of  $\langle A, \lesssim_1, \circ_1 \rangle$  if and only if  $A \subseteq B$ ,  $\lesssim_1 \subseteq \lesssim$ ,  $\circ_1 \subseteq \circ$ , and  $\langle B, \lesssim, \circ \rangle$  is a regularly dense ordered abelian group.

*Theorem 5.6.* Let  $\langle A, \lesssim_1, \circ_1 \rangle$  be a closed extensive structure. Then there is a

regularly dense ordered abelian group extension  $\langle B, \lesssim, \circ \rangle$  of  $\langle A, \lesssim_1, \circ_1 \rangle$  such that  $A \subseteq B^+$ .

*Proof.* Theorem 5.4 and Theorem 5.5.

*Theorem 5.7.* Let  $\langle A, \lesssim_1, \circ_1 \rangle$  be a closed extensive structure. Then there is a nonstandard model of the reals  $\langle *Re, *\mathcal{F} \rangle$  and a function  $f$  from  $A$  into  $*Re^+$  such that

- (1)  $x \prec_1 y$  iff  $f(x) < f(y)$ ; and
- (2)  $f(x \circ_1 y) = f(x) + f(y)$ .

*Proof.* By Theorem 5.6, let  $\langle B, \lesssim, \circ \rangle$  be a regularly dense ordered abelian group extension of  $\langle A, \lesssim_1, \circ_1 \rangle$ . Since  $A \subseteq B^+$ , by Theorem 4.8, let  $\langle *Re, *\mathcal{F} \rangle$  and  $f$  be such that  $f$  is an imbedding of  $A$  into  $*Re^+$ . Then  $f$  has the required properties (1) and (2).

*Definition 5.5.* Let  $\langle A, \lesssim, \circ \rangle$  be a closed extensive structure and  $x, y \in A$ . Let  $1x = x$  and for each  $n \in I^+$ ,  $(n + 1) = (nx) \circ x$ .  $x$  is said to be *commensurable with*  $y$  if and only if (1)  $x \lesssim y$  and for some  $n \in I^+$ ,  $nx \gtrsim y$ , or (2)  $y \lesssim x$  and for some  $n \in I^+$ ,  $ny \gtrsim x$ . As before, ‘is commensurable with’ can be shown to be an equivalence relation. A function  $s$  from  $A$  into  $Re^+$  is said to be a *closed extensive scale for*  $\langle A, \lesssim, \circ \rangle$  if and only if the following four conditions hold for all  $x, y, z$  in  $A$ :

- (1) if  $x$  is commensurable with  $y$  then  $s(x \circ y) = s(x) + s(y)$ ;
- (2) if  $x$  is commensurable with  $y$  then:
  - (i) if  $x \lesssim y$  then  $s(x) \leq s(y)$ , and
  - (ii) if  $s(x) < s(y)$  then  $x \prec y$ ;
- (3) if  $x \prec y$  and  $x$  is not commensurable with  $y$ , then  $s(x \circ y) = s(y)$ ; and
- (4) if  $x$  and  $y$  are commensurable,  $s(x) = s(y)$ , and  $x \circ z \lesssim y$ , then  $z$  is not commensurable with  $x$ .

*Theorem 5.8.* Let  $\langle A, \lesssim, \circ \rangle$  be a closed extensive structure. Then there is a closed extensive scale for  $\langle A, \lesssim, \circ \rangle$ .

*Proof.* Theorem 5.6 and Theorem 4.10.

*Theorem 5.9.* Let  $\langle A, \lesssim, \circ \rangle$  be a closed extensive structure,  $s$  and  $t$  be closed extensive scales for  $\langle A, \lesssim, \circ \rangle$ , and  $a \in A$ . Let  $T = \{x \in A \mid x \text{ is commensurable with } a\}$ . Then there is a positive real number  $r$  such that for each  $x \in T$ ,  $s(x) = rt(x)$ .

*Proof.* Theorem 5.6 and Theorem 4.11.

*Definition 5.6.* Let  $\langle A, \lesssim, \circ \rangle$  be an extensive structure,  $B$  be the domain of  $\circ$  and  $x \in A$ . Then, by definition  $1x = x$ . Furthermore, if  $nx$  has been defined for  $n \in I^+$  and  $(nx, x) \in B$  then, by definition,  $(n + 1)x = (nx) \circ x$ . If  $(nx, x) \notin B$  then  $(n + 1)x$  is not defined.

*Definition 5.7.* Let  $\langle A, \lesssim, \circ \rangle$  be an extensive structure. An element  $x \in A$  is said to be *small* if and only if for some  $y \in A$ ,  $mx \prec y$  for each  $m \in I^+$ .



*Theorem 5.10.* Let  $\langle A, \lesssim, \circ \rangle$  be an extensive structure. Suppose  $x \lesssim y$  and  $y$  is small. Then for each  $n \in I^+$ ,  $nx$  is small.

*Proof.* Let  $B$  be the domain of  $\circ$ . We will show by induction that  $nx$  is defined for each  $n \in I^+$  and that  $nx \lesssim ny$ . By definition,  $1x$  is defined and  $1x \lesssim 1y$ . Let  $p \in I^+$  and suppose that  $px$  has been defined and that  $px \lesssim py$ . Since  $(py, y) \in B$  and  $py \gtrsim px$ , by (3) of *Definition 5.1*,  $(y, px) \in B$  and  $(p + 1)y = (py) \circ y \gtrsim y \circ (px)$ . Since  $(y, px) \in B$  and  $y \gtrsim x$ , by (3) of *Definition 5.1*,  $(px, x) \in B$  and  $y \circ (px) \gtrsim (px) \circ x = (p + 1)x$ . Thus we have shown that  $(p + 1)x$  is defined and  $(p + 1)x \lesssim (p + 1)y$ . Thus by induction, for each  $n \in I^+$ ,  $nx \lesssim ny$ . Since for some  $z \in A$ ,  $ny < z$  for each  $n \in I^+$ ,  $nx < z$  for each  $n \in I^+$ . That is,  $x$  is small. Let  $m \in I^+$  and  $nx < z$  for each  $n \in I^+$ . Then  $n(mx) = (nm)x < z$  for each  $n \in I^+$ . That is,  $mx$  is small for each  $m \in I^+$ .

*Theorem 5.11.* Let  $\langle A, \lesssim, \circ \rangle$  be an extensive structure and  $D = \{x \mid x \in A \text{ and } x \text{ is small}\}$ . Suppose that  $D \neq \phi$ . Let  $\lesssim_1$  and  $\circ_1$  be the restrictions of  $\lesssim$  and  $\circ$  to  $D$ . Then  $\langle D, \lesssim_1, \circ_1 \rangle$  is a closed extensive structure.

*Proof.* We need only show that  $\circ_1$  is an operation on  $D$  since all the other conditions for a closed extensive structure follow immediately from the fact that  $\langle A, \lesssim, \circ \rangle$  is an extensive structure and from the definitions of  $\lesssim_1$  and  $\circ_1$ . Let  $B$  be the domain of  $\circ$  and let  $x, y$  be elements of  $D$ . To show that  $\circ_1$  is an operation on  $D$  we need only show that  $(x, y) \in B$  and  $x \circ y \in D$ .

Case 1. Suppose that  $x \lesssim y$ . Since  $(y, y) \in B$  and  $x \lesssim y$ , we have by condition (3) of *Definition 5.1* that  $(x, y) \in B$  and  $x \circ y \lesssim y \circ y$ . Since  $y \circ y \in D$ , by *Theorem 5.10*,  $x \circ y \in D$ .

Case 2. Suppose that  $x \succ y$ . Then by Case 1  $(y, x) \in B$ . Since  $y \lesssim y$ , by Condition (3) of *Definition 5.1*,  $(x, y) \in B$ . Since  $x \circ y \lesssim x \circ x$  and  $x \circ x \in D$ , by *Theorem 5.10*,  $x \circ y \in D$ .

*Definition 5.8.* Let  $\langle A, \lesssim, \circ \rangle$  be an extensive structure and  $x \in A$ . Then  $x$  is said to be *large* if and only if  $x$  is not small.

*Theorem 5.12.* Let  $\langle A, \lesssim, \circ \rangle$  be an extensive structure,  $x, y \in A$ , and  $x$  be small and  $y$  be large. Then  $x < y$ .

*Proof.* Since  $\lesssim$  is a weak ordering, either  $x < y$  or  $y \lesssim x$ . Since  $y$  is large, by *Theorem 5.10*, it cannot be the case that  $y \lesssim x$ . Therefore  $x < y$ .

In order to assure that an extensive structure can be represented, it is necessary to add conditions on the large elements to assure that they can be "measured." This is done in the following definition.

*Definition 5.9.* An extensive structure  $\langle A, \lesssim, \circ \rangle$  is said to be *good* if and only if for all large elements  $a, b$  of  $A$  the following two conditions hold:

(1) if  $a < b$  and for all small  $\alpha$ ,  $a \circ \alpha < b$ , then for some large  $c$  in  $A$ ,  $a \circ c \lesssim b$ ; and

(2) there are large  $c$  and  $d$  in  $A$  such that  $a < c \circ d$ .

*Definition 5.10.* Let  $\langle A, \lesssim, \circ \rangle$  be a good extensive structure and  $D$  be the set of large elements of  $A$ . Then the function  $f$  from  $D$  into  $Re^+$  is said to be an *extensive*

*imbedding of  $D$  into  $Re^+$*  if and only if the following three conditions hold for all  $x, y \in A$ :

- (1) if  $(x, y)$  is the domain of  $\circ$  then  $f(x \circ y) = f(x) + f(y)$ ;
- (2)  $f(x) < f(y)$  iff for some  $z \in D$ ,  $x \circ z \lesssim y$ ; and
- (3) if  $x \lesssim y$  then:  $f(x) = f(y)$  iff either  $x \sim y$  or for some small  $\alpha$  in  $A$ ,  $x \circ \alpha \gtrsim y$ .

*Theorem 5.13.* Let  $\langle A, \lesssim, \circ \rangle$  be a good extensive structure and  $D$  be the set of large elements of  $A$ . Then there is an extensive imbedding of  $D$  into  $Re^+$ .

*Outline of proof.* By Zorn’s lemma, let  $D'$  be a maximal subset of  $D$  such that if  $x, y \in D'$  and  $x < y$  then for some large  $z \in D$ ,  $x \circ z \lesssim y$ . Define the relation  $E$  on  $D$  as follows:  $xEy$  if and only if  $x \in D', y \in D$  and  $[(x < y$  and for some small  $\alpha \in A$ ,  $x \circ \alpha \gtrsim y)$  or  $(x \sim y)$  or  $(y < x$  and for some small  $\alpha \in A$ ,  $y \circ \alpha \gtrsim x)]$ . Then it is easy to show that for all  $y \in D$  there is an  $x \in D'$  such that  $xEy$ . Define  $\circ_1$  and  $\lesssim_1$  on  $D'$  as follows:  $x \circ_1 y = z$  iff there are  $x_1, y_1, z_1$  in  $D$  such that  $xEx_1, yEy_1, zEz_1$ , and  $x_1 \circ y_1 = z_1$ ; and  $x \lesssim_1 y$  iff for some  $x_1, y_1 \in D$ ,  $xEx_1, yEy_1$ , and  $x_1 \lesssim y_1$ . Then one can show that  $\langle D', \lesssim_1, \circ_1 \rangle$  is an Archimedean extensive structure. By *Theorem 5.1*, let  $g$  be an imbedding of  $\langle D', \lesssim_1, \circ_1 \rangle$  into  $Re^+$ . Define the function  $f$  from  $D$  into  $Re^+$  as follows: if  $y \in D$ , let  $f(y) = g(x)$  where  $x$  is such that  $xEy$ . Then it is easy to show that  $f$  is an extensive imbedding of  $D$  into  $Re^+$ .

We will now extend the definition of ‘commeasureability’ and ‘scale’ to extensive structures.

*Definition 5.11.* Let  $\langle A, \lesssim, \circ \rangle$  be an extensive structure and  $a, b \in A$ . Then  $a$  is said to be *extensively commensurable with  $b$*  if and only if (i)  $a$  and  $b$  are large elements of  $A$ , or (ii)  $a$  and  $b$  are small elements of  $A$  and  $a$  and  $b$  are commensurable (as defined for closed extensive structures—*Definition 5.5*). A function  $s$  from  $A$  into  $Re^+$  is said to be an *extensive scale* for  $\langle A, \lesssim, \circ \rangle$  if and only if the following four conditions hold for all  $x, y, z$  in  $A$ :

- (1) if  $x$  is extensively commensurable with  $y$  and  $(x, y) \in \text{domain } \circ$ , then  $s(x \circ y) = s(x) + s(y)$ ;
- (2) if  $x$  is extensively commensurable with  $y$  then: (i) if  $x \lesssim y$  then  $s(x) \leq s(y)$ , and (ii) if  $s(x) < s(y)$  then  $x < y$ ;
- (3) if  $x < y$  and  $x$  is not commensurable with  $y$ , then  $s(x \circ y) = s(y)$ ; and
- (4) if  $x$  and  $y$  are commensurable,  $s(x) = s(y)$ , and  $x \circ z \lesssim y$ , then  $z$  is not commensurable with  $x$ .

*Theorem 5.14.* Let  $\langle A, \lesssim, \circ \rangle$  be an extensive structure,  $s$  an extensive scale for  $\langle A, \lesssim, \circ \rangle$ ,  $a \in A$ , and  $T = \{x \in A \mid x \text{ is extensively commensurable with } a\}$ . Let  $r \in Re^+$  and  $t$  be a function from  $A$  into  $Re^+$  such that for all  $x \in A - T$ ,  $t(x) = s(x)$ , and for all  $x \in T$ ,  $t(x) = rs(x)$ . Then  $t$  is an extensive scale for  $\langle A, \lesssim, \circ \rangle$ .

*Proof.* Immediate from *Definition 5.11*.

*Theorem 5.15.* Let  $\langle A, \lesssim, \circ \rangle$  be a good extensive structure. Then there is an extensive scale  $s$  for  $\langle A, \lesssim, \circ \rangle$ . Furthermore, if  $t$  is another extensive scale for  $\langle A, \lesssim, \circ \rangle$ ,  $x \in A$ , and  $T = \{y \in A \mid y \text{ is extensively commensurable with } x\}$ , then there is a  $r \in Re^+$  such that for all  $z \in T$ ,  $s(z) = rt(z)$ .

*Proof.* Let  $S$  be the set of small elements of  $A$  and  $D$  be the set of large elements of  $A$ . If  $S = \phi$ , then  $\langle A, \lesssim, \circ \rangle$  is an Archimedean extensive structure and the theorem follows from *Theorem 5.1*. If  $D = \phi$ , then the theorem follows from *Theorems 5.8* and *5.9*. Therefore, assume that  $S \neq \phi$  and  $D \neq \phi$ . By *Theorem 5.8* let  $s_1$  be a closed extensive scale for  $S$  and by *Theorem 5.13*  $s_2$  be an extensive imbedding of  $D$  into  $Re^+$ . Let  $s = s_1 \cup s_2$ . Then it is easy to verify that  $s$  is an extensive scale for  $\langle A, \lesssim, \circ \rangle$ . It immediately follows from the definition of  $s$  and from *Theorems 5.9*, *5.1*, and the proof of *Theorem 5.13* that if  $x \in A$  and  $T = \{y \in A \mid y \text{ is extensively commensurable with } x\}$  and  $t$  is an extensive scale for  $\langle A, \lesssim, \circ \rangle$  then for some  $r \in Re^+$ ,  $s(z) = rt(z)$  for all  $z \in T$ .

The relationships between extensive structures and imbeddings into nonstandard models of the reals will be described in the next three theorems. The proofs of these theorems will be omitted.

*Theorems 5.16.* Let  $\langle A, \lesssim, \circ \rangle$  be a good extensive structure. Then there is a nonstandard model of the reals  $\langle *Re, *\mathcal{F} \rangle$  and a function  $f$  such that  $f$  is an imbedding of  $A$  into  $*Re^+$ .

The real number system is a *universal* measuring system in the sense that each Archimedean extensive structure can be imbedded into  $Re^+$ . So far we have only shown that for each extensive structure there is a nonstandard model of the reals into which it can be imbedded. We have not shown, for example, that two extensive structures can be imbedded in the same nonstandard model of the reals. By an intelligent application of the compactness theorem (*Theorem 1.1*), the following theorem can be proved:

*Theorem 5.17.* Let  $\mathcal{C}$  be a nonempty class of extensive structures. Then there is a nonstandard model of the reals  $\langle *Re, *\mathcal{F} \rangle$  such that each member of  $\mathcal{C}$  can be imbedded in  $*Re^+$ .

An even stronger version of *Theorem 3.4* can be proved by using *saturated models*. (See [1], Chapter 11.)

*Theorem 5.18.* Let  $\aleph$  be the cardinality of  $Re$  (alternatively,  $\aleph$  be a regular uncountable cardinal) and  $\mathcal{C}$  a nonempty class of extensive structures such that each member of  $\mathcal{C}$  has cardinality  $\leq \aleph$ . Assume the continuum hypothesis (alternatively, the generalized continuum hypothesis). Then there is a nonstandard model of the reals,  $\langle *Re, *\mathcal{F} \rangle$ , such that  $*Re$  has cardinality  $\aleph$  and each member of  $\mathcal{C}$  is imbeddable in  $*Re^+$ .

**6. Historical Note.** Axiomatic approaches to Archimedean extensive attributes were made by Helmholtz [2] and Hölder [3]. The axioms for Archimedean closed extensive structures are like Robert's and Luce's in [10]. The axioms for Archimedean extensive structures which are due to Krantz, *et. al.* [4], are a modification of axioms given in Luce and Marley [5]. Abraham Robinson has applied the compactness theorem to a wide variety of algebraic problems in Robinson [11], and has used

the nonstandard reals for the solution of many problems in Robinson [12]. Representation and uniqueness theorems for non-Archimedean additive conjoint structures and non-Archimedean qualitative probability structures are given in Narens [8], and non-Archimedean expected utility structures in Narens [9].

## REFERENCES

- [1] Bell, J. L. and Slomson, S. B. *Models and Ultraproducts*. Amsterdam: North-Holland Publishing Co., 1966.
- [2] Helmholtz, H. *Zählen und Messen Erkenntnis-Theoretisch Betrachtet. Philosophische Aufsätze Eduard Zeller gewidmet*. Leipzig, 1887. (English translation by C. L. Bryan, *Counting and measuring*. Princeton: Van Nostrand, 1930.)
- [3] Hölder, O. "Die Axiome der Quantität und die Lehre vom Mass." *Ber. Verh. Kgl. Sächsis Ges. Wiss. Leipzig, Math.-Phys. Classe* 53 (1901): 1-64.
- [4] Krantz, D. H., Luce, R. D., Suppes, P., and Tversky, A. *Foundations of Measurement*. Vol. I. New York: Academic Press, 1971.
- [5] Luce, R. D. and Marley, A. A. J. "Extensive Measurement when Concatenation is Restricted and Maximal Elements may Exist." In *Philosophy, Science, and Method: Essays in Honor of Ernest Nagel*. Edited by S. Morgenbesser, P. Suppes, and M. G. White. New York: St. Martin's Press, 1969. Pages 235-249.
- [6] Narens, L. "Why Tall Pygamas are Short." In *Social Science Working Papers #13*, School of Social Sciences, University of California, Irvine, November, 1972.
- [7] Narens, L. "Field Embeddings of Generalized Metric Spaces." In *Victoria Symposium on Nonstandard Analysis*. Edited by A. Hurd and P. Loeb. Heidelberg: Springer-Verlag, 1974.
- [8] Narens, L. "Minimal Conditions for Additive Conjoint Measurement and Qualitative Probability." (To appear in *Journal of Mathematical Psychology*, 1974.)
- [9] Narens, L. "Utility-Uncertainty Trade-off Structures." In *Social Sciences Working Papers #59*, School of Social Sciences, University of California, Irvine, August, 1974.
- [10] Roberts, F. S. and Luce, R. D. "Axiomatic Thermodynamics and Extensive Measurement." *Synthese* 18 (1968): 311-326.
- [11] Robinson, A. *Introduction to Model Theory and the Metamathematics of Algebra*. Amsterdam: North-Holland Publishing Co., 1965.
- [12] Robinson, A. *Non-Standard Analysis*. Amsterdam: North-Holland Publishing Co., 1966.
- [13] Robinson, A. and Zakon, E. "Elementary Properties of Ordered Abelian Groups." In *Transactions of the American Mathematical Society* 96 (1960): 222-236.