


## Summary: Neuroplasticity after Perinatal Brain Damage

- For both language and spatial cognition
   Perinatal brain damage results in developmental delay followed by subtle, long-term deficits
- Implications
  - Although some brain systems have a high level of genetic predisposition and thus suffer long-term harm when disrupted
  - Brain plasticity also exists, at least early in development, so that the cognitive functions that suffer early damage develop in an alternative manner.

Psych 9B -- PSB 11B

```
Development 4
```

17




Someone you know has had a major stroke. After 3 months of rehabilitation he/she still cannot walk, talk comprehensibly, or feed and take care of him/herself. At this point, you are advised by doctors that no further improvement is likely. The patient is discouraged by the his/her lack of improvement. Is it time to stop therapy and put the patient in a care facility?
A. Yes
B. No
C. I don't know

Psych 9B - PSB 11B

Development 4

19



	What Differentiates Patients with				
Better Recovery from Stroke					
	1. Use it or lose it	Failure to drive specific brain functions can lead to functional degradation.	s		
	2. Use it and improve it	Training that drives a specific brain function can enhance that function.			
	3. Specificity	The nature of the training experience dictates the nature of the plasticity.			
	<ol> <li>Repetition matters</li> </ol>	Induction of plasticity requires sufficie directed practice.	nt		
	Source: Kleim & Jones,	2008			
	Development 4	Psych 9B PSB 11B	21		

	What Differentiates Patients with Better Recovery from Stroke				
	5. Intensity matters	Induction of plasticity requires sufficient training intensity.	t		
	6. Timing matters	Different forms of plasticity occur at different times during training.			
	7. Rewards matter	The training experience must be sufficiently salient to induce plasticity.			
	8. Age matters	Training-induced plasticity occurs more readily in younger brains.	;		
-			-		
	Development 4	Psych 9B PSB 11B 2	22		

## Summary: Innate Brain Structures, Neuroplasticity, and Remapping

- Localization emerges during development
- It results from genetically determined processes
- The processes (and the resulting localization) are not unchangeable
  - □ They can be altered by environmental factors, experience, and damage
- Cortical remapping does happen

Development 4

- Although possible even in older adults
   It is more prevalent in younger brains
  - Psych 9B -- PSB 11B

23

